

Inhibition of *Enterococcus faecalis* by Calcium Peroxide

Yong Liang SU¹, Xiao Yan WANG²

Objective: To investigate the inhibition of *Enterococcus faecalis* by calcium peroxide (CaO_2).

Methods: The minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) of *Enterococcus faecalis* by CaO_2 and calcium hydroxide (Ca(OH)_2) were determined by direct exposure tests ($n = 10$). The inhibition zone of *E. faecalis* mycoderm treated with CaO_2 and Ca(OH)_2 paste (53% w/w) was observed using agar diffusion tests ($n = 20$). The inhibition of *E. faecalis* biofilms by CaO_2 /phosphate-buffered saline (PBS) and Ca(OH)_2 /PBS suspensions were observed using confocal laser scanning microscopy and the percentages of live bacteria in the biofilms calculated.

Results: The MIC of Ca(OH)_2 (4.5 to 5.5 mg/ml) was higher than the MIC of CaO_2 (2.0 to 2.5 mg/ml) ($P < 0.05$), and the MBC of Ca(OH)_2 (14.5 to 15.5 mg/ml) was higher than that of CaO_2 (3.0 to 3.5 mg/ml) ($P < 0.05$). No inhibition zone was observed for Ca(OH)_2 in agar diffusion tests, while the diameter of the inhibition zone around CaO_2 was 8.6 ± 0.4 mm. There were significant differences between groups in the percentages of surviving bacteria in *E. faecalis* biofilms after treatment ($P < 0.05$): group $\text{CaO}_2 < \text{group Ca(OH)}_2 < \text{group PBS} < \text{group BHI}$.

Conclusion: The inhibition of *E. faecalis* by CaO_2 was greater than that by Ca(OH)_2 .

Key words: biofilm, calcium hydroxide, calcium peroxide, *Enterococcus faecalis*
Chin J Dent Res 2016;19(2):109–113; doi: 10.3290/j.cjdr.a36181

The main objective of endodontic therapy is to eliminate bacteria from the root canal and to prevent the regrowth of residual microorganisms¹. Infected tissue removal is achieved by thorough biomechanical cleaning, including shaping and irrigating the canal with proteolytic disinfecting solution. However, thorough irrigation of root canals with antimicrobial solution might not be sufficient to eliminate all microorganisms from the root canal². Inability to completely eradicate microor-

ganisms and/or provide a bacterial seal can result in the failure of root canal therapy. Hence, the use of intracanal medicaments has been advocated to further reduce the number of microorganisms, for example, in teeth with chronic sinus infection, and in teeth that are undergoing revision of failed root canal procedures.

Enterococcus faecalis is a Gram-positive bacterium often isolated from persistent root canal infections. It can penetrate deeply into dentinal tubules and resist bactericidal substances commonly used in endodontic procedures^{3,4}. Its prevalence in asymptomatic, persistent endodontic infection is about 77%⁵. Siren et al⁶ found that *E. faecalis* was the most common enteric bacterium isolated from the root canal in both primary treatment and retreatment groups, and it appears as a monoinfection. *E. faecalis* has the capacity to live in dentinal tubules enduring prolonged periods of starvation^{7,8}, and *E. faecalis* biofilms have greater magnitude more resistant to antimicrobials than planktonic bacteria^{9,10}.

Ca(OH)_2 was first introduced to dentistry in 1920 and has been widely accepted as an intracanal medica-

1 First Clinical Division, Peking University School and Hospital of Stomatology, National Engineering Laboratory for Digital and Material Technology of Stomatology, Beijing Key Laboratory of Digital Stomatology, Beijing, P.R. China.

2 Department of Cariology and Endodontology, Peking University School and Hospital of Stomatology, National Engineering Laboratory for Digital and Material Technology of Stomatology, Beijing Key Laboratory of Digital Stomatology, Beijing, P.R. China.

Corresponding author: Dr Xiao Yan Wang, Department of Cariology and Endodontology, Peking University School and Hospital of Stomatology, 22# Zhongguancun South Avenue, HaiDian District, Beijing 100081, P.R. China. Tel: 86-10-82195525; Fax: 86-10-82195525; Email: wangxiaoyan@pkuss.bjmu.edu.cn

ment because of its antimicrobial properties, especially its action on Gram-negative bacteria^{11,12}. However, some authors have suspected it is effective in eliminating *E. faecalis*^{13,14}. The search for a better alternative has led to the use of newer antimicrobial agents such as CaO_2 .

CaO_2 is a white or yellowish solid peroxide, which can slowly decompose to release oxygen at a 'controlled' rate when in contact with hydrous media. Besides its stable oxygen releasing capability, CaO_2 also possesses capacities for bleaching, disinfection and deodorising¹⁵. Hence, CaO_2 has been widely used in agriculture, aquaculture and medicine¹⁵. In its natural state, CaO_2 dissolves in water to form hydrogen peroxide (H_2O_2) and Ca(OH)_2 and thus is considered a 'solid form' of H_2O_2 ¹⁵. The antibacterial properties of Ca(OH)_2 stems from its ability to increase the pH of a solution¹⁶. However, Mura et al presumed that reactive oxygen species (ROS) are released from CaO_2 and suggested that the species of ROS were H_2O_2 and the superoxide anion ($\bullet\text{O}_2^-$)¹⁷. Thus, high pH and/or ROS may be the sterilisation mechanism of CaO_2 , but no research has been published on this mechanism.

As a bactericidal alkali and oxygen-releasing agent, CaO_2 may be effective in controlling *E. faecalis* infections in the root canal. The purpose of this study was to evaluate the antibacterial effects of CaO_2 through a direct exposure method, agar diffusion method and by confocal laser scanning microscopy.

Materials and methods

Bacterial strain and medications

For this *in vitro* study, a single standard strain of *Enterococcus faecalis* (ATCC 29212; Lab of Microbiology, Beijing Stomatological Hospital, Capital Medical University) was selected. The strain was incubated on brain heart infusion broth (BHI) agar solid culture medium for 24 h. Single colonies were taken and incubated into BHI liquid medium (5% CO_2 at 37°C). After 24 h, the microbial cells were resuspended in BHI to achieve a final concentration of 3×10^8 cells/ml. Analytical grade Ca(OH)_2 and CaO_2 (China National Pharmaceutical Group Chemical Reagent, Shanghai, China) were used in this study.

Determination of MIC and MBC

BHI broth (4 ml) was placed in a centrifuge tube, then Ca(OH)_2 or CaO_2 was added. The concentrations of

Ca(OH)_2 were 1.5, 2.0, 2.5, 3.0, 3.5, 4.0, 4.5, 5.0, 5.5, 6.0, 14.0, 14.5, 15.0 and 15.5 mg/ml, respectively. The concentrations of CaO_2 were 1.5, 2.0, 2.5, 3.0, 3.5 and 4.0 mg/ml, respectively. Then, *E. faecalis* was added into the tubes (0.1 ml of 3×10^8 cells/ml). After 24 h, the MIC and MBC of Ca(OH)_2 and CaO_2 was determined. The MIC was defined as the minimum concentration without bacterial growth by macroscopic observation (without turbidity in the BHI broth). The MBC was defined as the minimum concentration where there was no bacterial growth on BHI agar plates. Samples were repeated ten times for each group. All data were analysed using SPSS 16.0 software (SPSS, Illinois, USA).

Observation of inhibition zone

Enterococcus faecalis suspensions (0.1 ml) were spread on BHI agar and incubated (5% CO_2 at 37°C). After the *E. faecalis* mycoderm formed on the BHI agar, CaO_2 paste (mixed with distilled water; 53% w/w) or Ca(OH)_2 paste (mixed with distilled water; 53% w/w) were put on the mycoderm. After cultivation for 24 h (5% CO_2 at 37°C), the inhibition zones were observed and measured ($n = 20$). All data were analysed using SPSS 16.0 software.

Observation of biofilm inhibition

Hydroxyapatite sheets (Clarkson Chromatography Company, Pennsylvania, USA) were placed in 24-well culture plates and immersed in 2 ml of BHI culture medium containing *E. faecalis* (3×10^8 cells/ml) for biofilm formation. The BHI culture medium containing *E. faecalis* was replaced every 2 days. After 14 days, the hydroxyapatite sheets were washed twice in phosphate-buffered saline (PBS) and divided into four groups: (i) BHI group: the hydroxyapatite sheets were immersed in 2 ml BHI medium and incubated for 48 h (5% CO_2 at 37°C). (ii) PBS group: the hydroxyapatite sheets were immersed in 2 ml PBS and incubated for 48 h. (iii) Ca(OH)_2 group: the hydroxyapatite sheets were immersed in 2 ml PBS containing 30 mg Ca(OH)_2 and incubated for 48 h. (iv) CaO_2 group: the hydroxyapatite sheets were immersed in 2 ml PBS containing 30 mg CaO_2 and incubated for 48 h. The hydroxyapatite sheets with biofilm were stained with LIVE/DEAD reagent (SYT09: PI: PBS = 1.5 μl : 1.5 μl : 1.0 ml) for 30 min. Then they were washed twice in PBS, treated with 1.5 μl anti-fluorescent quencher and observed after 24 h. All samples were examined under a confocal laser scanning microscope (Leica TCS-SPE; Leica Microsystems GmbH, Mannheim, Germany) at 63 \times magnification.

The viable, live bacteria were green, while the dead were red, and overlapping alive and dead bacteria produced either orange or yellow samples. Each sample was scanned from the outside of the biofilm (away from the hydroxyapatite sheet) to the inside along the Z-axis. The biofilm observed was divided into three layers: outer one-third, middle one-third and inner one-third. The percentage of viable cells was calculated for each layer using IpWin32 software and the data was analysed using SPSS 16.0.

Results

Table 1 showed that the MIC of CaO_2 towards *E. faecalis* was 2.0 to 2.5 mg/ml, while the MIC of Ca(OH)_2 was 4.5 to 5.5 mg/ml. The MBC of CaO_2 was 3.0 to 3.5 mg/ml, while that of Ca(OH)_2 was 14.5 to 15.5 mg/ml. There were statistically significant differences between the two treatments ($P < 0.05$).

Figure 1 showed the pH of CaO_2 and Ca(OH)_2 solutions before and after incubation with *E. faecalis*. The pH increased along with an increasing concentration of Ca(OH)_2 . When the concentration of Ca(OH)_2 was 5 mg/ml, the pH reached 11.87. However, the pH ranged from 8.10 to 9.50 in a solution of CaO_2 (1.5 to 5.0 mg/ml). After incubation with bacteria for 24 h, the pH of the Ca(OH)_2 solution decreased. A much slighter decrease in pH was observed for the CaO_2 solution.

Figure 2 showed typical inhibition zones of *E. faecalis* by Ca(OH)_2 and CaO_2 on BHI agar plates. No inhibition zone was observed for Ca(OH)_2 , while the inhibition zone of CaO_2 was 8.6 ± 0.4 mm in diameter.

Figures 3 and 4 showed the results of biofilm inhibition. The percentage of live bacteria in the inner layer of the biofilms was higher than that in the outer layer for all groups. Ca(OH)_2 killed some *E. faecalis* in biofilms (Fig 3). CaO_2 treatment resulted in the lowest percentage of live bacteria in the whole of the biofilm, as well as in each of the inner, middle and outer layers. The percentage of live bacteria in the CaO_2 group < Ca(OH)_2 group < PBS group < BHI group ($P < 0.05$).

Discussion

CaO_2 showed a lower MIC and MBC towards *E. faecalis* than Ca(OH)_2 . CaO_2 forms H_2O_2 and Ca(OH)_2 when it dissolves in water and is thus considered a 'solid form' of H_2O_2 ¹⁵. The sterilisation mechanism of Ca(OH)_2 mainly depends on a high pH value caused by dissociation of hydroxyl ions; the high pH may destroy cell membranes, denature structural proteins and enzymes, and damage DNA¹⁸. However, in this study, the pH of CaO_2 solution

Table 1 MIC and MBC of CaO_2 and Ca(OH)_2 towards *Enterococcus faecalis*.

	MIC (mg/ml)	MBC (mg/ml)
CaO_2	2.0-2.5	3.0-3.5
Ca(OH)_2	4.5-5.5	14.5-15.5




Fig 1 The pH of Ca(OH)_2 and CaO_2 suspensions of different concentrations before and after bacteriostatic action. CH indicates Ca(OH)_2 , CP indicates CaO_2 , 24 h means that pHs were measured after incubation of the sample with *Enterococcus faecalis* for 24 h; pHs of other samples were measured after 0 h.




Fig 2 Inhibition zones of *Enterococcus faecalis* by Ca(OH)_2 and CaO_2 paste (53% w/w) on BHI agar plates after cultivation for 24 h. CH indicates Ca(OH)_2 ; CP indicates CaO_2 ; red arrows indicate inhibition zones.




Fig 3 Biofilm images obtained by confocal laser scanning microscopy followed by 3D reconstruction; with a 63x magnification. Green indicates live bacteria; red indicates dead bacteria. BHI - brain heart infusion; PBS - phosphate-buffered saline; CH - Ca(OH)₂; CP - CaO₂.

was around 8.10 to 9.50, which is much lower than the antibacterial pH (around 12)¹⁹. This indicates that the antibacterial mechanism of CaO₂ may arise from the oxygen free radicals released from hydrogen peroxide. Block et al confirmed that hydrogen peroxide generated oxygen free radicals and the reaction with macromolecules (such as membrane lipids and DNA) lead to bacterial death²⁰. However, a previous study showed that 3% H₂O₂ solution, which also generated oxygen free radicals, had no significant inhibitory effect on planktonic *E. faecalis*²¹. In that study, there was not enough contact time (with the method of irrigating immediately) for H₂O₂ to kill the bacteria; the short lifetime of the oxygen

free radicals generated by hydrogen peroxide led to a low bacterial inhibition effect. However, in the present study, the CaO₂ was in contact with the *E. faecalis* for 24 h, and produced oxygen free radicals continuously via a slow interaction with water. Thus we were able to observe an obvious inhibitory effect on planktonic *E. faecalis*.

In the agar diffusion test used for observation of inhibition zones, the test results depend on the solubility and diffusivity of medicaments in the agar, rather than just their efficacy against the organism^{22,23}. In this study, there was no inhibition zone of *E. faecalis* around Ca(OH)₂ paste, while there was a clear inhibition zone around CaO₂ paste. This indicates that the solubility and diffusivity of CaO₂ in agar appears to be better than that of Ca(OH)₂.

A previous study reported that *E. faecalis* biofilms grew well on the hydroxyapatite sheet and the biofilm entered a relatively stable state after 14 days, after which the total number of bacteria did not change significantly²⁴. Therefore, we selected the hydroxyapatite sheet as a biofilm substrate in this study. Based on the MIC, MBC and inhibition zone results, it was not a surprise that CaO₂ showed stronger inhibition effects towards *E. faecalis* biofilms than Ca(OH)₂. However, it should be noted that the *E. faecalis* biofilm was not completely killed by CaO₂ in this study, indicating that the biofilm has a stronger capacity to resist killing than the planktonic state. The extracellular matrix (a gel polymer) of biofilms has a buffer action against antibacterial agents²⁵. In addition, as bacteria in the inner layer




Fig 4 The percentage of viable cells in different layers of *Enterococcus faecalis* biofilms treated with antimicrobials: CH - Ca(OH)₂; CP - CaO₂; BHI - brain heart infusion; PBS - phosphate-buffered saline.

relatively lack nutrition, their starvation state means they could be resistant to antibacterial agents.

Within the limitations of this study, CaO₂ showed greater inhibitory effects against planktonic and biofilm forms of *Enterococcus faecalis* than Ca(OH)₂.

Acknowledgments

The authors gratefully acknowledged the Lab of Microbiology, Peking University School and Hospital of Stomatology.

Conflicts of interest

The authors reported no conflicts of interest related to this study.

Author contributions

- Dr Yong Liang Su for carrying out the research and preparing the paper.
- Dr Xiao Yan Wang for the design of the study and directing the research.

References

1. Aravind V, Gopikrishna D, Kandaswamy, Rajan K Jeyavel. Comparative evaluation of the antimicrobial efficacy of five endodontic root canal sealers against *Enterococcus faecalis* and *Candida albicans*. *J Conserv Dent* 2006;9:2–12.
2. Bystrom A, Claesson R, Sundqvist G. The antibacterial effect of camphorated paramonochlorophenol, camphorated phenol and calcium hydroxide in the treatment of infected root canals. *Endod Dent Traumatol* 1985;1:170–175.
3. Williams JM, Trope M, Caplan DJ, Shugars DC. Detection and quantitation of *E. faecalis* by real-time PCR (qPCR), reverse transcription-PCR (RT-PCR), and cultivation during endodontic treatment. *J Endod* 2006;32:715–721.
4. Haapasalo M, Orstavik D. In vitro infection and disinfection of dental tubules. *J Dent Res* 1987;66:1375–1379.
5. Siqueira JF Jr, Rôças IN. Polymerase chain reaction-based analysis of microorganisms associated with failed endodontic treatment. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2004;97:85–94.
6. Siren EK, Haapasalo MP, Ranta K, Salmi P, Kerosuo EN. Microbiological findings and clinical treatment procedures in endodontic cases selected for microbiological investigation. *Int Endod J* 1997;30:91–95.
7. Love RM. *Enterococcus faecalis*—a mechanism for its role in endodontic failure. *Int Endod J* 2001;34:399–405.
8. Figdor D, Davies JK, Sundqvist G. Starvation survival, growth and recovery of *Enterococcus faecalis* in human serum. *Oral Microbiol Immunol* 2003;18:234–239.
9. Distel JW, Hatton JF, Gillespie MJ. Biofilm formation in medicated root canals. *J Endod* 2002;28:689–693.
10. Costerton JW, Stewart PS, Greenberg EP. Bacterial biofilms: a common cause of persistent infections. *Science* 1999;284:1318–1322.
11. Athanassiadis B, Abbott PV, Walsh LJ. The use of calcium hydroxide, antibiotics and biocides as antimicrobial medicaments in endodontics. *Aust Dent J* 2007;52(Suppl 1): S64–S82.
12. Menezes MM, Valera MC, Jorge AO, Koga-Ito CY, Camargo CH, Mancini MN. In vitro evaluation of the effectiveness of irrigants and intracanal medicaments on microorganisms within root canals. *Int Endod J* 2004;37:311–319.
13. Lana PE, Scelza MF, Silva LE, Mattos-Guaraldi AL, Hirata Júnior R. Antimicrobial activity of calcium hydroxide pastes on *Enterococcus faecalis* cultivated in root canal systems. *Braz Dent J* 2009;20:32–36.
14. Chai WL, Hamimah H, Cheng SC, Sallam AA, Abdullah M. Susceptibility of *Enterococcus faecalis* biofilm to antibiotics and calcium hydroxide. *J Oral Sci* 2007;49:161–166.
15. Ma Y, Zhang BT, Zhao L, Guo G, Lin JM. Study on the generation mechanism of reactive oxygen species on calcium peroxide by chemiluminescence and UV-visible spectra. *Luminescence* 2007;22:575–580.
16. Siqueira JF Jr, Lopes HP. Mechanisms of antimicrobial activity of calcium hydroxide: a critical review. *Int Endod J* 1999;32:361–369.
17. Mura A, Medda R, Longu S, Floris G, Rinaldi AC, Padiglia A. A Ca²⁺/calmodulin-binding peroxidase from *Euphorbia latex*: novel aspects of calcium-hydrogen peroxide cross-talk in the regulation of plant defenses. *Biochemistry* 2005;44:14120–14130.
18. Estrela C, Rodrigues de Araújo Estrela C, Bammann LL, Pecora JD. Two methods to evaluate the antimicrobial action of calcium hydroxide paste. *J Endod* 2001;27:720–723.
19. Behnen MJ, West LA, Liewehr FR, Buxton TB, McPherson JC 3rd. Antimicrobial activity of several calcium hydroxide preparations in root canal dentin. *J Endod* 2001;27:765–767.
20. Block SS. Peroxygen compounds. *Disinfection, Sterilization and Preservation*, ed 4. Philadelphia: Lea & Febiger, 1991:167–181.
21. Wang DM, Gao XJ, Shen S. Comparison of antimicrobial efficacy of four endodontic irrigants using an in vitro model infected by *Enterococcus faecalis* [In Chinese]. *Zhonghua Kou Qiang Yi Xue Za Zhi* 2007;42:223–224.
22. Basrani B, Tjäderhane L, Santos JM, Pascon E, Grad H, Lawrence HP, Friedman S. Efficacy of chlorhexidine- and calcium hydroxide-containing medicaments against *Enterococcus faecalis* in vitro. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2003;96:618–624.
23. Gomes BP, Vianna ME, Sena NT, Zaia AA, Ferraz CC, de Souza Filho FJ. In vitro evaluation of the antimicrobial activity of calcium hydroxide combined with chlorhexidine gel used as intracanal medicament. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2006;102:544–550.
24. Guerreiro-Tanomaru JM, de Faria-Júnior NB, Duarte MA, Ordinola-Zapata R, Graeff MS, Tanomaru-Filho M. Comparative analysis of *Enterococcus faecalis* biofilm formation on different substrates. *J Endod* 2013;39:346–350.
25. Basrani B, Santos JM, Tjäderhane L, Grad H, Gorduysus O, Huang J, Lawrence HP, Friedman S. Substantive antimicrobial activity in chlorhexidine-treated human root dentin. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2002;94:240–245.